

Counting Models of Genus One Curves

Mohammad Sadek

Abstract

Let C be a soluble smooth genus one curve over a Henselian discrete valuation field. There is a unique minimal Weierstrass equation defining C up to isomorphism. In this paper we consider genus one equations of degree n defining C , namely a (generalised) binary quartic when $n = 2$, a ternary cubic when $n = 3$, and a pair of quaternary quadrics when $n = 4$. In general, minimal genus one equations of degree n are not unique up to isomorphism. We explain how the number of these equations varies according to the Kodaira symbol of the Jacobian of C . Then we count these equations up to isomorphism over a number field of class number 1.

1 Introduction

Let E be an elliptic curve defined over a number field K . It is known that $E(K)$ is a finitely generated abelian group. If we manage to find $E(K)/nE(K)$ for any integer $n \geq 2$, then a finite amount of computation will determine $E(K)$. The method of n -descent is one of the methods which enable us to get a bound on $E(K)/nE(K)$, $n \geq 2$. Indeed, the n -descent computes the n -Selmer group of E which contains $E(K)/nE(K)$ as a subgroup. The difference between the two groups is the n -torsion of the Tate-Shafarevich group.

An element of the n -Selmer group can be represented as a geometric object, namely as an everywhere locally soluble curve defined by a *genus one equation of degree n* given as follows:

If $n = 1$, a Weierstrass equation

$$y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6. \quad (1)$$

Two genus one equations of degree 1 with coefficients in a Dedekind domain R are *R -equivalent* if they are related by the substitutions: $x' = u^2x + r$ and $y' = u^3y + su^2x + t$,

where $r, s, t \in R$, $u \in R^*$. The group of transformations $[u; r, s, t]$ is called $\mathcal{G}_1(R)$. We set $\det([u; r, s, t]) = u^{-1}$.

If $n = 2$, a (generalised) binary quartic

$$y^2 + (\alpha_0 x^2 + \alpha_1 xz + \alpha_2 z^2)y = ax^4 + bx^3z + cx^2z^2 + dxz^3 + ez^4. \quad (2)$$

Two genus one equations of degree 2 with coefficients in R are *R-equivalent* if they are related by the substitutions: $x' = m_{11}x + m_{21}z$, $z' = m_{12}x + m_{22}z$ and $y' = \mu^{-1}y + r_0x^2 + r_1xz + r_2z^2$, where $\mu \in R^*$, $r_i \in R$ and $M = (m_{ij}) \in \text{GL}_2(R)$. The group of transformations $[\mu, (r_i), M]$ is called $\mathcal{G}_2(R)$. We set $\det([\mu, (r_i), M]) = \mu \det M$.

If $n = 3$, a ternary cubic

$$F(x, y, z) = ax^3 + by^3 + cz^3 + a_2x^2y + a_3x^2z + b_1y^2x + b_3y^2z + c_1z^2x + c_2z^2y + mxyz = 0. \quad (3)$$

Two genus one equations of degree 3 with coefficients in R are *R-equivalent* if they are related by multiplying by $\mu \in R^*$, then substituting $x' = m_{11}x + m_{21}y + m_{31}z$, $y' = m_{12}x + m_{22}y + m_{32}z$ and $z' = m_{13}x + m_{23}y + m_{33}z$, where $M = (m_{ij}) \in \text{GL}_3(R)$. The group of transformations $[\mu, M]$ is called $\mathcal{G}_3(R)$. Set $\det([\mu, M]) = \mu \det M$.

If $n = 4$, then C is described by 2 quadratic forms in 4 variables. Two genus one equations of degree 4 with coefficients in R are *R-equivalent* if they are related by the substitutions: $F'_i = m_{i1}F_1 + m_{i2}F_2$ for $M = (m_{ij}) \in \text{GL}_2(R)$, and then $x'_j = \sum_{i=1}^4 n_{ij}x_i$ for $N = (n_{ij}) \in \text{GL}_4(R)$. The group of transformations $[M, N]$ is called $\mathcal{G}_4(R)$. Set $\det([M, N]) = \det M \det N$.

For $n \leq 4$, we associate invariants $c_{4,\phi}$, $c_{6,\phi}$ and *discriminant* Δ_ϕ to a genus one equation ϕ of degree n where $\Delta_\phi = (c_{4,\phi}^3 - c_{6,\phi}^2)/1728$. Moreover, ϕ defines a smooth curve of genus one if and only if $\Delta_\phi \neq 0$. The invariants $c_{4,\phi}$, $c_{6,\phi}$ and Δ_ϕ are of weights $r = 4, 6$ and 12 respectively, in other words if $F \in \{c_{4,\phi}, c_{6,\phi}, \Delta_\phi\}$, then $F \circ g = (\det g)^r F$ for all $g \in \mathcal{G}_n$. For more details on these genus one equations and their classical invariants see [1]. We scale the above invariants according to [7] and [5].

For using n -descent to search for points on an elliptic curve E , we need the coefficients of the genus one equations described above to be small. That can be achieved by *Reduction* and *Minimisation*. By reducing genus one equations, we mean reducing the size of the coefficients by a unimodular linear change of coordinates, which does not change the invariants. To minimise genus one equations, we need to make the associated invariants smaller. For references see [2] and [15] for $n = 2$, [6] for $n = 3$, and [16] for $n = 4$. An algorithmic approach to the minimisation problem can be found in [5]. In this paper we will be concerned with the minimisation question.

We work throughout over a perfect Henselian discrete valuation field K with algebraic closure $\overline{K}$ and ring of integers $\mathcal{O}_K$. We fix a uniformiser t , a normalisation ν and write $k = \mathcal{O}_K/t\mathcal{O}_K$. Set $S = \text{Spec } \mathcal{O}_K$.

Definition 1.1. A genus one equation ϕ of degree n , $n \leq 4$, with $\Delta_\phi \neq 0$ is

- (a) *integral* if the defining polynomials have coefficients in $\mathcal{O}_K$.
- (b) *minimal* if it is integral and $\nu(\Delta_\phi)$ is minimal among all the valuations of the discriminants of the integral genus one equations K -equivalent to ϕ .

It has been shown in [5] and [12] that if C is a soluble smooth curve defined by a genus one equation ϕ of degree n , then the minimal discriminant associated to ϕ has the same valuation as the minimal discriminant associated to the Jacobian elliptic curve.

The S -scheme $\mathcal{C}$ defined by an integral genus one equation $\phi : y^2 + g(x, z)y = f(x, z)$ of degree 2 is the scheme obtained by glueing $\{y^2 + g(x, 1)y = f(x, 1)\} \subset \mathbb{A}_S^2$ and $\{v^2 + g(1, u)v = f(1, u)\} \subset \mathbb{A}_S^2$ via $x = 1/u$ and $y = x^2v$. It comes with a natural morphism $\mathcal{C} \rightarrow \mathbb{P}_S^1$ given on these affine pieces by $(x, y) \mapsto (x : 1)$ and $(u, v) \mapsto (1 : u)$. The S -scheme defined by an integral genus one equation ϕ of degree n , $n = 1$ or $n \geq 3$, is the subscheme $\mathcal{C} \subset \mathbb{P}_S^m$ defined by ϕ , where $m = 2$ when $n = 1$, and $m = n - 1$ when $n \geq 3$. In [12], we gave the following criterion for minimality.

Theorem 1.2. *Let ϕ be an integral genus one equation of degree n , $n \leq 4$. Assume moreover that ϕ defines a normal S -scheme $\mathcal{C}$, and that $\mathcal{C}_K$ is smooth with $\mathcal{C}_K(K) \neq \emptyset$. Then ϕ is minimal if and only if the minimal desingularisation of $\mathcal{C}$ is isomorphic to the minimal proper regular model of the Jacobian of $\mathcal{C}_K$.*

Unlike elliptic curves, the minimisations of integral genus one equations of degree n are not unique under the action of the group $\mathcal{G}_n(\mathcal{O}_K)$. In this paper we study the non-uniqueness of minimisation. In order to achieve this goal we generalise the scheme-theoretic terminology of minimal Weierstrass models for elliptic curves to *minimal degree- n -models* for genus one curves.

Definition 1.3. Let ϕ be a genus one equation of degree n . Let $C \rightarrow \mathbb{P}_K^m$ be the genus one curve over K defined by ϕ , where $m = 2$ when $n = 1$ and $m = n - 1$ otherwise. A *(minimal) degree- n -model* for $C \rightarrow \mathbb{P}_K^m$ is a pair $(\mathcal{C}, \alpha)$ where $\mathcal{C} \subset \mathbb{P}_S^m$ is an S -scheme defined by a (minimal) integral genus one equation of degree n , and $\alpha : \mathcal{C}_K \cong C$ is an isomorphism defined by a K -equivalence of genus one equations of degree n , i.e., α is defined by an element of $\mathcal{G}_n(K)$. If there is no confusion, then we will omit mentioning α in $(\mathcal{C}, \alpha)$ and write $\mathcal{C}$ instead.

An isomorphism $\beta : (\mathcal{C}_1, \alpha_1) \cong (\mathcal{C}_2, \alpha_2)$ of degree- n -models is an isomorphism $\beta : \mathcal{C}_1 \cong \mathcal{C}_2$ of S -schemes defined by an $\mathcal{O}_K$ -equivalence of genus one equations of degree n , i.e., β is defined by an element of $\mathcal{G}_n(\mathcal{O}_K)$, with $\beta_K = \alpha_2^{-1}\alpha_1$.

Now we show why we insist on remembering the isomorphism α in our definition of a degree- n -model. In the case $C(K) \neq \emptyset$, we pick $P \in C(K)$. Then we identify (C, P) with the Jacobian elliptic curve E . The automorphism group $\text{Aut}(C)$ of C fits in an exact sequence

$$0 \rightarrow E \rightarrow \text{Aut}(C) \rightarrow \text{Aut}(E, 0) \rightarrow 0.$$

The first map is $Q \mapsto \tau_Q$, where τ_Q is the translation by Q . Let H be a hyperplane section on C . We are interested in automorphisms λ of C such that $\lambda^*H \sim H$. But $\tau_Q^*H \sim H$ if and only if $nQ = 0$. Hence the elements of $\text{PGL}_n(K)$ that act fixed-point-free on C correspond precisely to $E[n](K)$. Therefore, we can have more than one K -isomorphism between $\mathcal{C}_K$ and C defined by elements in $\mathcal{G}_n(K)$.

Liu proved that there is a bijection between minimal degree-2-models for $C \rightarrow \mathbb{P}_K^1$, up to isomorphism, and the multiplicity-1 irreducible components in the special fiber of the quotient of the minimal proper regular model by a hyperelliptic involution, see ([8], Corollaire 5).

In §3, we perform explicit computations on the minimal proper regular model of an elliptic curve. In §4, these computations are used to count minimal degree- n -models for a soluble smooth genus one curve $C \rightarrow \mathbb{P}_K^{n-1}$, $n \in \{2, 3, 4\}$, up to isomorphism. We relate the number of minimal degree- n -models for $C \rightarrow \mathbb{P}_K^{n-1}$, up to isomorphism, to the cardinality of a finite set which depends only on the Kodaira Symbol of the Jacobian of C and on the divisor defining the morphism $C \rightarrow \mathbb{P}_K^{n-1}$, see Theorem 4.2. Then, using strong approximation, we prove that the number of minimal global degree- n -models for $C \rightarrow \mathbb{P}_K^{n-1}$ over a number field K of class number 1 is the product of minimal degree- n -models for $C \rightarrow \mathbb{P}_{K_\nu}^{n-1}$, where K_ν is the completion of K at a finite place ν and ν runs over a finite set of bad primes of the Jacobian of C .

We have to mention that over a Henselian discrete valuation field, our counting results work for any residue characteristic if the Jacobian has multiplicative reduction, while for additive reduction we assume $\text{char } k \neq 2$ when $n = 2$, and $\text{char } k \neq 2, 3$ when $n = 3, 4$. It seems to us that these conditions on the residue characteristic for the additive reduction case are not necessary, but we have not been able to eliminate them.

2 Isomorphisms of degree- n -models

Let C be a smooth genus one curve over K . Assume that $C(K) \neq \emptyset$. Let E be the Jacobian of C with minimal proper regular model E^{min} . We define an S -model for C to be an integral, projective, flat, normal S -scheme $f : X \rightarrow S$ of dimension 2 such that its generic fiber $\mathcal{C}_K$ is isomorphic to C . The special fiber of $\mathcal{C}$ will be called $\mathcal{C}_k$.

Let $P \in C(K)$. Fix an isomorphism $\beta : C \rightarrow E$ such that $\beta(P) = 0_E$. The isomorphism β identifies (C, P) with $(E, 0_E)$. We observe that β extends to an isomorphism between the minimal proper regular model of C and E^{min} . In [12] we proved that a minimal degree- n -model $\mathcal{C}$ for a smooth genus one curve $C \rightarrow \mathbb{P}_K^{n-1}$ is normal, and hence is an S -model for C . Moreover, we proved that the minimal desingularisation of $\mathcal{C}$ is isomorphic to E^{min} , see Theorem 1.2. Furthermore, we have

$$\mathcal{C} = \text{Proj}\left(\bigoplus_{m=0}^{\infty} H^0(E^{min}, \mathcal{O}_{E^{min}}(mD))\right),$$

where D is a divisor on E^{min} which intersects the irreducible components of the strict transform of $\mathcal{C}_k$ in E_k^{min} and intersects no other components in E_k^{min} . The divisor $D|_E$ is linearly equivalent to the hyperplane section divisor defining C .

Let Γ be an irreducible component of $\mathcal{C}_k$. By the *type* of Γ we mean the ordered pair $(\text{mult}_k(\Gamma), \deg_k(\Gamma))$, where $\text{mult}_k(\Gamma)$ and $\deg_k(\Gamma)$ are the multiplicity and degree of Γ respectively.

Theorem 2.1. *Let C be a smooth genus one curve with $C(K) \neq \emptyset$. Let $(\mathcal{C}_1, \alpha_1)$ and $(\mathcal{C}_2, \alpha_2)$ be two minimal degree- n -models for $C \rightarrow \mathbb{P}_K^{n-1}$, $n = 2, 3, 4$. Set $\alpha = \alpha_2^{-1}\alpha_1$ and denote its extension $\mathcal{C}_1 \dashrightarrow \mathcal{C}_2$ by $\tilde{\alpha}$. Then the following statements are equivalent.*

- (i) $(\mathcal{C}_1, \alpha_1)$ and $(\mathcal{C}_2, \alpha_2)$ are isomorphic as degree- n -models for $C \rightarrow \mathbb{P}_K^{n-1}$.
- (ii) For every irreducible component Γ of $(\mathcal{C}_2)_k$, $\tilde{\alpha}^*\Gamma$ is an irreducible component of $(\mathcal{C}_1)_k$ with the same type as Γ .

PROOF: (i) implies (ii): Let $\alpha := \alpha_2^{-1}\alpha_1 : (\mathcal{C}_1)_K \rightarrow (\mathcal{C}_2)_K$. By the definition of isomorphic degree- n -models, the map α extends to an S -isomorphism $\tilde{\alpha} : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ which is defined by an element in $\mathcal{G}_n(\mathcal{O}_K)$. Therefore, if Γ is an irreducible component of $(\mathcal{C}_2)_k$, then $\tilde{\alpha}^*\Gamma$ is an irreducible component of $(\mathcal{C}_1)_k$ with the same type as Γ .

(ii) $\Rightarrow$ (i): We want to show that $\tilde{\alpha}$ is defined by an element in $\mathcal{G}_n(\mathcal{O}_K)$. Since the minimal proper regular models of $(\mathcal{C}_1)_K$ and $(\mathcal{C}_2)_K$ are identified with E^{min} , statement (ii) implies that both $(\mathcal{C}_1)_k$ and $(\mathcal{C}_2)_k$ have the same strict transform in E_k^{min} . Let D_i be a defining divisor of $\mathcal{C}_i$ as a contraction in E^{min} , i.e.,

$$\mathcal{C}_i = \text{Proj}\left(\bigoplus_{m \geq 0} H^0(E^{min}, \mathcal{O}_{E^{min}}(mD_i))\right).$$

Set $\mathcal{L}_i = \mathcal{O}_{E^{min}}(D_i)$ and $\mathcal{L}_{i,k} = \mathcal{L}_i|_{E_k^{min}}$. Let Γ be an irreducible component of $(\mathcal{C}_2)_k$. Consider the strict transform $\tilde{\Gamma}$ of the irreducible components Γ and $\tilde{\alpha}^*\Gamma$ in E_k^{min} . Since Γ and $\tilde{\alpha}^*\Gamma$ have the same type, it follows that $\deg_k \mathcal{L}_{1,k}|_{\tilde{\Gamma}} = \deg_k \mathcal{L}_{2,k}|_{\tilde{\Gamma}}$. For any

irreducible component Λ which is not a strict transform of a component of $(\mathcal{C}_i)_k$, we have $\deg_k \mathcal{L}_{i,k}|_\Lambda = 0$. Now since each irreducible component of E_k^{min} is isomorphic to $\mathbb{P}_k^1$, we have $D_1|_{E_k^{min}}$ is linearly equivalent to $D_2|_{E_k^{min}}$ as they have the same degree on every irreducible component of E_k^{min} . It follows that $\mathcal{L}_{1,k} \cong \mathcal{L}_{2,k}$.

We are given that $C(K) \neq \emptyset$, $\mathcal{L}_1|_{(C_1)_K} \cong \mathcal{L}_2|_{(C_2)_K}$ and we established the isomorphism $\mathcal{L}_{1,k} \cong \mathcal{L}_{2,k}$. These conditions imply that $\mathcal{L}_1 \cong \mathcal{L}_2$, see ([9], Exercise 9.1.13 (b)). Therefore, $H^0(E^{min}, \mathcal{L}_1)$ and $H^0(E^{min}, \mathcal{L}_2)$ are isomorphic as $\mathcal{O}_K$ -modules, and $\tilde{\alpha}$ is a change of basis of a free $\mathcal{O}_K$ -module of rank n . Hence $\tilde{\alpha}$ is defined by an element in $\mathcal{G}_n(\mathcal{O}_K)$. $\square$

Corollary 2.2. *Let C be a smooth genus one curve. Assume that $C(K) \neq \emptyset$. Assume moreover that the Jacobian E of C has either reduction types I_0 or I_1 . Then there is a unique minimal degree- n -model for $C \rightarrow \mathbb{P}_K^{n-1}$.*

3 Computing in E^{min}

The following lemma will play an essential rule in the way we construct divisors on minimal proper regular models.

Proposition 3.1. *Let C/K be a smooth genus one curve with $C(K) \neq \emptyset$. Let E be the Jacobian of C with minimal proper regular model E^{min} . Fix a closed point $x \in E_k^{min}$ such that x lies on one and only one irreducible component Γ of E_k^{min} , of multiplicity $r \geq 1$. Then there exists a point $P \in C$ such that $\overline{\{P\}} \cap E_k^{min} = \{x\}$ and $[K(P) : K] = r$, where $\overline{\{P\}}$ is the Zariski closure of $\{P\}$ in E^{min} .*

PROOF: See ([9], Exercise 9.2.11 (c)). $\square$

It is easy to check that if L is a totally ramified extension of K with $[L : K] = m$, $m = 2, 3, 4$, then the maximal unramified extension L^{un} of L is a Galois extension of the maximal unramified extension K^{un} of K with $L^{un} = K^{un}(\sqrt[m]{t})$.

Let E be an elliptic curve over K , with minimal proper regular model E^{min} . Let $E^0(K)$ be the group of rational points of E with non-singular reduction. In this section we define a map $\delta_m : \Phi_K^m(E) \rightarrow \Phi_K(E)$, where $\Phi_K^m(E)$ is the set of multiplicity- m irreducible components of E_k^{min} , and $\Phi_K(E) := \Phi_K^1(E)$ is the group of components $E(K)/E^0(K)$. In the remainder of this section we assume that the residue field k is algebraically closed and hence every finite extension of K is a totally ramified Galois extension, and the reduction type of E is split.

3.1 The component group

The component group $\Phi_K(E) = E(K)/E^0(K)$ is finite. More precisely, if E has multiplicative reduction, then $E(K)/E^0(K)$ is a cyclic group of order $\nu(\Delta)$; otherwise, $E(K)/E^0(K)$ has order 1, 2, 3, or 4, see ([14], Chapter IV, Corollary 9.2).

If E has reduction I_n , $n \geq 0$, then E_k^{min} consists of an n -gon. There exists an isomorphism $\delta_1 : \Phi_K(E) \cong \mathbb{Z}/n\mathbb{Z}$ such that if we number the irreducible components in $\Phi_K(E)$ consecutively from 0 to $n-1$, then the image of the i th component in $\Phi_K(E)$ is $i \in \mathbb{Z}/n\mathbb{Z}$. If E has additive reduction, then we fix an isomorphism $\delta_1 : \Phi_K(E) \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ when E has reduction of type I_{2m}^* , $m \geq 0$, and $\delta_1 : \Phi_K(E) \cong \mathbb{Z}/n\mathbb{Z}$ for some $n \in \{1, 2, 3, 4\}$ for the other additive reduction types. From now on we will not distinguish between $\Phi_K(E)$ and its isomorphic image under δ_1 .

We note that if E has one of the reduction types I_n , $n \geq 0$, II, III, or IV, then E_k^{min} consists only of multiplicity-1 components, hence $\Phi_K^m(E) = \emptyset$ when $m \geq 2$.

We define the map $\delta_m : \Phi_K^m(E) \rightarrow \Phi_K(E)$, $m \geq 2$, as follows: Every irreducible component in $\Phi_K^m(E)$ is isomorphic to $\mathbb{P}_k^1$. Let x be a k -point of $\Theta \in \Phi_K^m(E)$ which lies on no other component. Let $P \in E(\overline{K})$ be a point which reduces to x such that $[K(P) : K] = m$, see Proposition 3.1. Let σ be a generator of $\text{Gal}(K(P)/K)$. The sum of the Galois orbit $P + \dots + P^{\sigma^{m-1}}$ of P is a point in $E(K)$. We set $\delta_m(\Theta)$ to be the image of this sum in $\Phi_K(E) = E(K)/E^0(K)$. It will be clear from the description of the map δ_m given below that $\delta_m(\Theta)$ does not depend on the choice of x nor on the choice of P .

Since $K(P)/K$ is a totally ramified extension, the Galois group $\text{Gal}(K(P)/K)$ is the inertia group of $K(P)/K$. In other words, $\text{Gal}(K(P)/K)$ fixes every irreducible component in E_k^{min} . Hence if $P \in E(\overline{K})$ lies above an irreducible component Γ of E_k^{min} , then every point in the Galois orbit of P lies above Γ as well.

3.2 Reduction type I_n^* , $n \geq 0$

Assume that E/K has reduction of type I_n^* , $n \geq 0$. The special fiber E_k^{min} contains a sequence of $(n+1)$ multiplicity-2 components and no components of higher multiplicities. Tate's algorithm, see ([14], Chapter IV, §9), can be used to write down explicit equations for the components in $\Phi_K^2(E)$, hence we can determine conditions for a point in E defined over $K(\sqrt{t})$ to lie above one of these multiplicity-2 components, and compute the sum of the Galois orbit of this point. We will write $a_{i,r}$ for $t^{-r}a_i$, and x_r, y_r for $t^{-r}x, t^{-r}y$ respectively.

The proof of Tate's algorithm shows that if E has additive reduction of one of the types I_n^* , $n \geq 0$, IV^* , III^* or II^* , then we can assume that there exists a minimal

Weierstrass equation of the form


$$y^2z + a_1xyz + a_3yz^2 = x^3 + a_2x^2z + a_4xz^2 + a_6z^3, \quad a_i \in \mathcal{O}_K,$$

for E/K , such that $t \mid a_1, a_2, t^2 \mid a_3, a_4$ and $t^3 \mid a_6$. Moreover, the identity component is given by $\Gamma_0 : z = 0$ and is attached to a multiplicity-2 component $V_{-1} : y_1^2 = 0$.

If E has reduction of type I_n^* , $n \geq 0$, then the non-identity multiplicity-1 component attached to V_{-1} is $\Gamma : x_1 + \tilde{a}_{2,1} = 0$. The other multiplicity-2 components of E^{min} are given by

$$V_l : \begin{cases} y_u^2 + \tilde{a}_{3,u}y_u - \tilde{a}_{6,2u} = 0 & \text{if } l = 2u - 3 \text{ is odd,} \\ \tilde{a}_{2,1}x_u^2 + \tilde{a}_{4,u+1}x_u + \tilde{a}_{6,2u+1} = 0 & \text{if } l = 2u - 2 \text{ is even.} \end{cases}$$

In addition, V_l consists of two distinct lines precisely when $l = n = \nu(\Delta) - 6$. The special fiber E_k^{min} is as in the following figure.



Let $K_2 = K(\sqrt{t})$ be the unique quadratic tame Galois extension of K . A point lying above a k -point on V_l , where $l = 2u - 2$, and on no other component is a point in $E(K_2)$ of the form $(\alpha t^{u+1} + \beta t^{u+1/2}, y)$, where $y \in K_2$, $t^{u+1} \mid y$, $\alpha \in \mathcal{O}_K$ and $\beta \in \mathcal{O}_K^*$. We have $\beta \in \mathcal{O}_K^*$, since otherwise this point would reduce to a point on V_{l+1} . Similarly, a point lying above a k -point on V_l , where $l = 2u - 3$, and on no other component is a point in $E(K_2)$ of the form $(x, \alpha t^{u+1} + \beta t^{u+1/2})$, where $x \in K_2$, $t^u \mid x$, $\alpha \in \mathcal{O}_K$ and $\beta \in \mathcal{O}_K^*$.

We recall the addition formula on elliptic curves. Let $P_1, P_2 \in E(\overline{K})$. If $P_1 \neq \pm P_2$, then

$$x(P_1 + P_2) = \lambda^2 + a_1\lambda - a_2 - x(P_1) - x(P_2), \quad \text{and} \quad y(P_1 + P_2) = -(\lambda + a_1)x(P_1 + P_2) - \mu - a_3,$$

where

$$\lambda = \frac{y(P_2) - y(P_1)}{x(P_2) - x(P_1)}, \quad \mu = \frac{y(P_1)x(P_2) - y(P_2)x(P_1)}{x(P_2) - x(P_1)}.$$

Lemma 3.2. *Assume E has one of the reduction types I_n^* , $n \geq 0$, IV^* , III^* , or II^* . Let $K_2 = K(\sqrt{t})$ with $\text{Gal}(K_2/K) = \langle \sigma \rangle$. Then a point $P \in E(K_2)$ lying above the component $y_1^2 = 0$ satisfies $P + P^\sigma \in E^0(K)$.*

PROOF: If $P^\sigma = -P$, then $P + P^\sigma = 0 \in E^0(K)$ and we are done. So, assume $x(P) \notin K$. Now $P = (\alpha't + \beta't^{3/2}, \alpha t^2 + \beta t^{3/2})$, where $\alpha', \beta' \in \mathcal{O}_K$, $\alpha \in \mathcal{O}_K$ and $\beta \in \mathcal{O}_K^*$. In the addition

formula given above, $\lambda = \beta/\beta' \in K$. Since $\nu(\beta') \geq 0$, it follows that $\nu(x(P + P^\sigma)) \leq 0$ because $t \mid a_1, a_2$. Hence the reduction of $P + P^\sigma$ is not a singular point, i.e., $P + P^\sigma \in E^0(K)$. $\square$

Proposition 3.3. *Assume that E/K has reduction type I_n^* , $n \geq 0$. Let $K_2 = K(\sqrt{t})$ with $\text{Gal}(K_2/K) = \langle \sigma \rangle$.*

- (i) *The multiplicity-1 component $\Gamma : x_1 + \tilde{a}_{2,1} = 0$ is of order 2 in $\Phi_K(E)$.*
- (ii) *Let $P \in E(K_2)$ be a point lying above V_l where $l = 2u - 3$. Then $P + P^\sigma \in E^0(K)$.*
- (iii) *Let $P \in E(K_2)$ be a point lying above V_l where $l = 2u - 2$. Then $P + P^\sigma$ reduces to a point on Γ .*

PROOF: (i) If P lies above $\Gamma : x_1 + \tilde{a}_{2,1} = 0$, then $-P$ lies above Γ as well. It follows that the inverse of Γ when considered as an element of $\Phi_K(E)$ is itself.

(ii) If $P^\sigma = -P$, then $P + P^\sigma = 0 \in E^0(K)$ and we are done. So, assume that $x(P) \notin K$. We can assume that $P = (\alpha't^u + \beta't^{u+1/2}, \alpha t^{u+1} + \beta t^{u+1/2})$, where $\alpha, \beta \in \mathcal{O}_K$ with $\nu(\beta) = 0$ and $\alpha', \beta' \in \mathcal{O}_K$. We have $\lambda = \beta/\beta'$, hence $\nu(\lambda) \leq 0$. Thus $\nu(x(P + P^\sigma)) \leq 0$, and $P + P^\sigma \in E^0(K)$.

(iii) We can assume that $P = (\alpha t^{u+1} + \beta t^{u+1/2}, y)$, where $y \in K_2$, $t^{u+1} \mid y$ and $\alpha, \beta \in \mathcal{O}_K$ with $\nu(\beta) = 0$. Therefore, we have $\nu(\lambda) > 0$ where $\lambda := (y(P) - y(P^\sigma))/(x(P) - x(P^\sigma))$. Following the addition formula we have $x(P + P^\sigma) = \lambda^2 + a_1\lambda - a_2 - x(P) - x(P^\sigma)$. Dividing by t and using that $t \mid a_1, a_2$, we get that $x_1(P + P^\sigma) = -\tilde{a}_{2,1} \pmod{t}$. $\square$

In Proposition 3.3, since $\Gamma : x_1 + \tilde{a}_{2,1} = 0$ has order 2, we have $\delta_1(\Gamma) = 2 \in \Phi_K(E) \cong \mathbb{Z}/4\mathbb{Z}$, when n is odd. When n is even, we know that $\delta_1 : \Phi_K(E) \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$, therefore every non-identity irreducible component has order 2. From now on, we will fix δ_1 such that $\delta_1(\Gamma) = (1, 1)$.

Corollary 3.4. *Assume E/K has reduction type I_n^* , $n \geq 0$. Let $\Phi_K^2(E) = \{V_{-1}, V_0, \dots, V_{n-1}\}$ be as above. Then $\delta_2 : \Phi_K^2(E) \rightarrow \Phi_K(E)$ is determined according to the following table.*

(n, l)	$(2m + 1, 2u - 3)$	$(2m + 1, 2u - 2)$	$(2m, 2u - 3)$	$(2m, 2u - 2)$
$\delta_2(V_l)$	0	2	(0, 0)	(1, 1)

3.3 Reduction types IV^* , III^* and II^*

For reduction types IV^* , III^* and II^* , we will not use Tate's algorithm to write down explicit equations for the components in $\Phi_K^2(E)$ because the defining equations of the components in $\Phi_K^m(E)$, $m \geq 2$, are complicated. Instead, we make use of our knowledge of the reduction types of E over totally ramified field extensions of K , and use the projection formula to exploit the symmetry of the graphs associated to E_k^{min} .

Lemma 3.5. *Let E/K be an elliptic curve. Let $K_m = K(t^{1/m})$. If E/K has reduction type IV^* , then E/K_3 has good reduction if $\text{char } k \neq 3$. If E/K has reduction III^* , then E/K_2 has reduction I_0^* and E/K_3 has reduction III .*

PROOF: Let $t_m = t^{1/m}$. We recall from ([14], Chapter IV, §9) that E/K has a minimal Weierstrass equation


$$y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$$

with $\nu(a_1) \geq 1, \nu(a_2) \geq 2, \nu(a_3) = 2, \nu(a_4) \geq 3, \nu(a_6) \geq 5$ if the reduction is IV^* , and $\nu(a_1) \geq 1, \nu(a_2) \geq 2, \nu(a_3) \geq 3, \nu(a_4) = 3, \nu(a_6) \geq 5$ if the reduction is III^* . Making substitutions $x \mapsto t_m^{2(m-1)}x$ and $y \mapsto t_m^{3(m-1)}y$, we will have a minimal Weierstrass equation for E/K_m of the form

$$y^2 + a_1t_m^{-(m-1)}xy + t_m^{-3(m-1)}a_3y = x^3 + a_2t_m^{-2(m-1)}x^2 + t_m^{-4(m-1)}a_4x + t_m^{-6(m-1)}a_6$$

with reduction type as specified in the statement of the lemma. $\square$

If E has one of the reduction types IV^* , III^* or II^* , then E_k^{min} is as in the figures below. Recall that the multiplicity-1 component Γ_i corresponds to $i \in \mathbb{Z}/n\mathbb{Z}$ under the isomorphism δ_1 . The numbers on the irreducible components refer to the multiplicities.



Proposition 3.6. *Let E/K be an elliptic curve with minimal proper regular model E^{min} . Let Ξ_m denote a component of multiplicity- m in E_k^{min} .*

(i) *If E/K has reduction IV^* and $\text{char } k \neq 3$, then*

Ξ_m	Θ_0	Θ_1	Θ_2	Λ_0
$\delta_m(\Xi_m)$	0	2	1	0

(ii) *If E/K has reduction III^* and $\text{char } k \neq 2, 3$, then*

Ξ_m	Θ_0	Θ_1	Θ_2	Λ_0	Λ_1	Ψ
$\delta_m(\Xi_m)$	0	0	1	0	1	0

(iii) *If E/K has reduction II^* , then $\delta_m(\Xi_m) = 0$ for every irreducible component Ξ_m in E_k^{min} .*

PROOF: Let $K_m = K(t^{1/m})$, $m \in \{2, 3, 4\}$. Let $\text{Gal}(K_m/K) = \langle \sigma_m \rangle$. We want to find $l \in \Phi_K(E)$ such that if $P \in E(K_m)$ reduces to a point on Ξ_m , then $\sum_{i=0}^{m-1} P^{\sigma_m^i}$ reduces to a point on Γ_l and therefore $\delta_m(\Xi_m) = l$. If $Q \in E(K)$, then we will denote the translation-by- Q automorphism by $\tau_Q : E \rightarrow E$. This K -automorphism extends to give an $\mathcal{O}_K$ -automorphism

$\tau'_Q : E^{min} \rightarrow E^{min}$, see e.g. ([14], Chapter IV, Proposition 4.6). In (i) and (ii), we know that $\Theta_0 : y_1^2 = 0$ and $\delta_2(\Theta_0) = 0$, see Lemma 3.2.

(i) Let $P \in E(K_2)$ be a point lying above Θ_i , $i = 1, 2$, and above no other component. Now let $Q \in E(K)$ be a point lying above Γ_{2i} . We have $\tau'_Q(\Gamma_i) = \Gamma_0$, and the projection formula implies that $\tau'_Q(\Gamma_i) \cdot \tau'_Q(\Theta_i) = \Gamma_i \cdot \Theta_i$, see ([9], Theorem 9.2.12), therefore $\tau'_Q(\Theta_i) = \Theta_0$. Hence $\tau_Q(P) + \tau_Q(P)^{\sigma_2} \in E^0(K)$, because $\delta_2(\Theta_0) = 0$, i.e., $(P + Q) + (P^{\sigma_2} + Q) \in E^0(K)$. In other words, $P + P^{\sigma_2} \in Q + E^0(K)$, i.e., $P + P^{\sigma_2}$ lies above Γ_{2i} .

Since E/K_3 has reduction type I_0 , see Lemma 3.5, $\bar{k} = k$ and $\text{char } k \neq 3$, then $E(K_3)$ is divisible by 3. For $P \in E(K_3)$ lying above Λ_0 , there is $Q \in E(K_3)$ with $P = 3Q$. Now $P + P^{\sigma_3} + P^{\sigma_3^2} = 3(Q + Q^{\sigma_3} + Q^{\sigma_3^2}) \in E^0(K)$ because $\Phi_E(K) \cong \mathbb{Z}/3\mathbb{Z}$.

(ii) Following the same argument as in (i) with τ_Q and $Q \in E(K) \setminus E^0(K)$, and remembering that $\delta_2(\Theta_0) = 0$, we have $\delta_2(\Theta_1) = 0$. Let $\mathcal{E}_m$ be the minimal proper regular model of E/K_m , $m = 2, 3$. Indeed, there is a morphism

$$\psi_m : \mathcal{E}_m \rightarrow \text{Norm}(E^{min} \times_{\mathcal{O}_K} \mathcal{O}_{K_m}) \rightarrow E^{min},$$

where $\text{Norm}(E^{min} \times_{\mathcal{O}_K} \mathcal{O}_{K_m})$ is the normalisation of $E^{min} \times_{\mathcal{O}_K} \mathcal{O}_{K_m}$, see for example ([9], §10.4) and ([10], pp. 10-11). The minimal proper regular model $\mathcal{E}_m$ is obtained from the latter normalisation by contracting (-1) -curves.

By virtue of Lemma 3.5, the reduction type of E/K_2 is I_0^* . There exists a minimal Weierstrass equation for E/K_2

$$y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6, \quad a_i \in \mathcal{O}_{K_2},$$

where $x^3 + a_2x^2 + a_4x + a_6 = (x - t\alpha_1)(x - t\alpha_2)(x - t\alpha_3)$, $\alpha_i \in \mathcal{O}_{K_2}$, such that $(\mathcal{E}_2)_k$ consists of $\Gamma_0 : z = 0$, $V_{-1} : y_1^2 = 0$ and the three distinct lines of $x_1^3 + \tilde{a}_{2,1}x_1^2 + \tilde{a}_{4,2}x_1 + \tilde{a}_{6,3} = 0$. Let $P_i = (t\alpha_i, 0)$. The reduction map $E(K_2)[2] = \{0, P_1, P_2, P_3\} \rightarrow \Phi_{K_2}(E)$ is surjective as each of these points lies on a different multiplicity-1 component. Now σ_2 fixes one of the α_i 's and swaps the other two. It follows that σ_2 fixes two multiplicity-1 components $\Gamma_{(0,0)}$ and Γ' of $(\mathcal{E}_2)_k$ and swaps the other two multiplicity-1 components Γ_* , $\Gamma_*^{\sigma_2}$. Note that $\Gamma_{(0,0)}$ and Γ' lie above Γ_0 and Γ_1 in E_k^{min} respectively. Since $\text{char } k \neq 2$ and $[K_2 : K]$ divides the multiplicities of Θ_2 and Ψ , it is known that Γ_* and $\Gamma_*^{\sigma_2}$ lie above Θ_2 under the morphism ψ_m , see for example ([9], Remark 10.4.8). Therefore, if P lies above Θ_2 in E^{min} , then it lies above either Γ_* or $\Gamma_*^{\sigma_2}$ in $\mathcal{E}_2$. Since $\Phi_{K_2}(E) \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$, we have $P + P^{\sigma_2}$ lies above Γ' in $(\mathcal{E}_2)_k$, and hence it lies above Γ_1 in E_k^{min} . Furthermore, the multiplicity-2 irreducible component V_{-1} of $(\mathcal{E}_2)_k$ lies above Ψ . We notice that $K_2 = K_4^G$ where $G = \{1, \sigma_4^2\}$. Therefore, if $P \in E(K_4)$ lies above Ψ , then $\sum_{i=0}^3 P^{\sigma_4^i} = (P + P^{\sigma_4}) + (P + P^{\sigma_4})^{\sigma_4^2} \in E^0(K_2)$, because $\delta_2(V_{-1}) = (0, 0)$, i.e., this sum lies above $\Gamma_{(0,0)}$ of $(\mathcal{E}_2)_k$ which lies above the identity component Γ_0 of E_k^{min} .

We know that E/K_3 has reduction type III, see Lemma 3.5. If $P \in E(K_3)$, then P lies on a multiplicity-1 component of $(\mathcal{E}_3)_k$. Moreover, since $\text{char } k \neq 3$, according to the description of the morphism ψ_3 given in ([10], pp. 10-11) if P lies on Λ_0 , then P lies on the identity

component of $(\mathcal{E}_3)_k$, and if P lies on Λ_1 , then P lies on the other multiplicity-1 component of $(\mathcal{E}_3)_k$. Thus we are done by observing that $\Phi_{K_3}(E) \cong \mathbb{Z}/2\mathbb{Z}$.

(iii) This is straightforward because $\sum_{i=0}^{m-1} P^{\sigma_i} \in E(K)$, and $E(K) = E^0(K)$ for reduction type II^* . $\square$

4 Counting minimal models

Let K be a Henselian discrete valuation field with ring of integers $\mathcal{O}_K$. We fix a uniformiser t . We denote the normalised valuation on K by ν . We assume that the residue field k is perfect with algebraic closure $\bar{k}$.

Let C be a smooth genus one curve over K such that $C(K) \neq \emptyset$. Let E be the Jacobian elliptic curve of C with minimal proper regular model $E^{\min}$. When E has additive reduction, we will assume moreover that $\text{char } k \neq 2$ for $n = 2$, and $\text{char } k \neq 2, 3$ for $n = 3, 4$.

Let $Q \in C(K)$. Again we fix an isomorphism $\beta : C \rightarrow E$ such that $\beta(Q) = 0_E$. So we can dispense with C and write E instead. If D is an effective K -rational divisor of degree n on E , then the Riemann-Roch theorem implies that there exists a point $P \in E(K)$ such that $D \sim (n-1).0_E + P$. Therefore, the equation defining $C \rightarrow \mathbb{P}_K^{n-1}$ is an equation for the double cover of the projective line $E \rightarrow \mathbb{P}_K^1$ given by the divisor class $[0_E + P]$ when $n = 2$, or the image of E when it is embedded in $\mathbb{P}_K^{n-1}$ by the divisor class $[(n-1).0_E + P]$ when $n = 3, 4$. Let ψ_E be the surjective homomorphism in the following short exact sequence

$$0 \rightarrow E^0(K) \rightarrow E(K) \xrightarrow{\psi_E} \Phi_E(K) \rightarrow 0.$$

Let $S^n(P)$, $n \leq 4$, be the set consisting of unordered tuples $(\Gamma_1, \dots, \Gamma_l)$, where Γ_i is an irreducible component of $E_{\bar{k}}^{\min}$, satisfying the following conditions:

- (i) $\sum_{i=1}^l \text{mult}_{\bar{k}}(\Gamma_i) = n$.
- (ii) $\sum_{i=1}^l \delta_{m_i}(\Gamma_i) = \psi_E(P)$, where $m_i = \text{mult}_{\bar{k}}(\Gamma_i)$.
- (iii) $(\Gamma_1^\sigma, \dots, \Gamma_l^\sigma) = (\Gamma_1, \dots, \Gamma_l)$ for every $\sigma \in \mathcal{G}_k$.

We define the map

$$\lambda_n : \{\text{minimal degree-}n\text{-models for } E \rightarrow \mathbb{P}_K^{n-1} \text{ up to isomorphism}\} \rightarrow S^n(P); (\mathcal{C}, \alpha) \mapsto (\Gamma_1, \dots, \Gamma_l),$$

such that

- (i) $(\Gamma_1, \dots, \Gamma_l)$ consists of the irreducible components of the strict transform of $\mathcal{C}_{\bar{k}}$ in $E_{\bar{k}}^{\min}$.
- (ii) If Γ is the strict transform of a component Γ' in $\mathcal{C}_{\bar{k}}$, then Γ appears in $(\Gamma_1, \dots, \Gamma_l)$ as many times as $\deg_{\bar{k}} \Gamma'$. In particular, we have $\sum_{i=1}^l \text{mult}_{\bar{k}}(\Gamma_i) = n$. See [11] and [4] for the complete classification of the multiplicities and degrees of the irreducible components of $\mathcal{C}_k$ when $n = 3, 4$. The case $n = 2$ is clear.

Lemma 4.1. *The map λ_n is well-defined.*

PROOF: To show that λ_n is well-defined we need to prove (i) if $\mathcal{C}$ and $\mathcal{C}'$ are minimal isomorphic degree- n -models for $E \rightarrow \mathbb{P}_K^{n-1}$, then $\lambda_n(\mathcal{C}) = \lambda_n(\mathcal{C}')$, and (ii) $\lambda_n(\mathcal{C}) \in S^n(P)$. To prove (i) we know that the special fibers of any two isomorphic degree- n -models for $E \rightarrow \mathbb{P}_K^{n-1}$ have the same irreducible components with the same types, therefore the corresponding tuples of both models are the same, see Theorem 2.1.

For (ii) we will show first that if $(\Gamma_1, \dots, \Gamma_l)$ is the tuple associated to $\mathcal{C}$, then $\sum_{i=1}^l \delta_{m_i}(\Gamma_i) = \psi_E(P)$ where $m_i = \text{mult}_k(\Gamma_i)$. Let D be a divisor on $E^{\min}$ such that $\mathcal{C}$ is obtained from $E^{\min}$ by contraction using D , see Theorem 1.2. The divisor D intersects the Γ_i 's and no other components. Endowing the Γ_j 's with the reduced structure, we have $D \cdot \Gamma_j = \deg_k \Gamma'_j$, where Γ'_j is an irreducible component of $\mathcal{C}_k$ whose strict transform in $E_k^{\min}$ is Γ_j . Let $(x_1, \dots, x_l)$ be a tuple of all intersection points of D with the irreducible components $(\Gamma_1, \dots, \Gamma_l)$ where $x_i \in \Gamma_i$. Note that x_i may be repeated in $(x_1, \dots, x_l)$ if Γ_i is the strict transform of a component whose degree is greater than 1. We have $D|_E \sim (n-1) \cdot 0 + P$.

Let $m_r = \max\{m_j : 1 \leq j \leq l\}$. It is clear that for Γ_j , $j = 1, \dots, l$, the multiplicity $m_j \in \{1, m_r\}$, since otherwise we will have $1 < m_j < m_r$ which implies that $m_r \geq 3$, hence $m_r + m_j \geq 5$ which contradicts that $\sum_{i=1}^l m_i = n \leq 4$. By virtue of Proposition 3.1, there exists a closed point $P_j \in E$ such that $[K(P_j) : K] = m_j$ and $\overline{\{P_j\}} \cap \Gamma_j = \{x_j\}$. Let $K_j = K(P_j)$ and $\text{Gal}(K_j^{\text{un}}/K^{\text{un}}) = \langle \sigma_j \rangle$. Let $\mathcal{E}$ be the minimal proper regular model of E/K_r^{un} . Since x_j lies on one and only one component of $E_k^{\min}$, there are exactly m_j points $\{y_1, \dots, y_{m_j}\}$ of $\mathcal{E}$ lying above x_j , and each of these points lies on a multiplicity-1 component of $\mathcal{E}_{\bar{k}}$, see ([9], Remark 10.4.8). Now we view D as a divisor on $\mathcal{E}$. For $Q \in E$, $\tilde{Q}$ will denote its reduction. We have

$$\tilde{P} = (\text{sum } D|_E)^{\sim} = \text{sum } D|_{\mathcal{E}_{\bar{k}}} = \sum_{j=1}^l \sum_{i=1}^{m_j} y_i = \sum_{j=1}^l \sum_{i=1}^{m_j} \tilde{P}_j^{\sigma_j^i}.$$

The second equality holds because D intersects $\mathcal{E}_{\bar{k}}$ only in multiplicity-1 components. Therefore, we have $P - \sum_{j=1}^l \sum_{i=1}^{m_j} \tilde{P}_j^{\sigma_j^i} \in E^0(K)$. Applying the surjective group homomorphism $\psi_E : E(K) \rightarrow \Phi_K(E)$, we get $\psi_E(P) = \sum_{j=1}^l \psi_E(\sum_{i=1}^{m_j} \tilde{P}_j^{\sigma_j^i}) = \sum_{j=1}^l \delta_{m_j}(\Gamma_j)$.

To show that $\lambda_n(\mathcal{C})$ is $\mathcal{G}_k$ -invariant, we know that D intersects the irreducible components of the strict transform of $\mathcal{C}_{\bar{k}}$ in $E_k^{\min}$ and no other components. If Γ is the strict transform of an irreducible component in $\mathcal{C}_{\bar{k}}$, then there is an $x \in (D|_{E_k^{\min}}) \cap \Gamma$. Moreover, $H^0(E^{\min}, \mathcal{O}_{E^{\min}}(mD))$ is a free $\mathcal{O}_K$ -module for every $m \geq 0$. Therefore, $D|_{E_K^{\min}}$ and $D|_{E_{\bar{k}}^{\min}}$ are K - and k -rational divisors respectively. Thus $x^{\sigma} \in (D|_{E_K^{\min}})^{\sigma} \cap \Gamma^{\sigma} = (D|_{E_{\bar{k}}^{\min}}) \cap \Gamma^{\sigma}$ for any $\sigma \in \mathcal{G}_k$. In particular, Γ^{σ} is the strict transform of an irreducible component of $\mathcal{C}_{\bar{k}}$. $\square$

We will show that λ_n is bijective, and so the set of isomorphism classes of minimal degree- n -models for $E \rightarrow \mathbb{P}_K^{n-1}$ is identified with $S^n(P)$. Consequently, we have the following theorem.

Theorem 4.2. *Let E/K be an elliptic curve and $P \in E(K)$. Let $E \rightarrow \mathbb{P}_K^{n-1}$ be the morphism determined by the divisor class $[(n-1).0_E + P]$, $n \in \{2, 3, 4\}$. Let $\delta_m : \Phi_K^m(E) \rightarrow \Phi_K(E)$ be the function defined in §3. Then there is a bijection between the set of minimal degree- n -models for $E \rightarrow \mathbb{P}_K^{n-1}$ up to isomorphism and the disjoint union of the following $\mathcal{G}_k$ -invariant sets.*

(i) *The set of unordered n -tuples*

$$S_1(n, P) = \{(a_1, \dots, a_n) : a_i \in \Phi_K(E) \mid a_1 + \dots + a_n = \psi_E(P)\}^{\mathcal{G}_k},$$

(ii) *the set $S_n(P) = \{a \in \Phi_K^n(E) \mid \delta_n(a) = \psi_E(P)\}^{\mathcal{G}_k}$,*

and

if $n \geq 3$

(iii) *the set $S_{(n-1,1)}(P) = \{(a, b) \in \Phi_K^{n-1}(E) \times \Phi_K(E) \mid \delta_{n-1}(a) + b = \psi_E(P)\}^{\mathcal{G}_k}$,*

and

if $n = 4$

(iv) *the set $S_{(2,1,1)}(P) = \{(a, b, c) \in \Phi_K^2(E) \times \Phi_K(E) \times \Phi_K(E) \mid \delta_2(a) + b + c = \psi_E(P)\}^{\mathcal{G}_k}$,
where the order of b and c is immaterial,*

(v) *the set of unordered pairs*

$$S_{(2,2)}(P) = \{(a, b) \in \Phi_K^2(E) \times \Phi_K^2(E) \mid \delta_2(a) + \delta_2(b) = \psi_E(P)\}^{\mathcal{G}_k}.$$

We will denote the number of multiplicity-1 irreducible components of $E_{\bar{k}}^{min}$ which are defined over k by c_k , this is the *Tamagawa number* of E/K . Now we get the following corollary as a direct consequence of Theorem 4.2.

Corollary 4.3. *Let E/K be an elliptic curve and let $P \in E(K)$. Let $E \rightarrow \mathbb{P}_K^{n-1}$ be the morphism determined by the divisor class $[(n-1).0_E + P]$. The number of minimal degree- n -models for $E \rightarrow \mathbb{P}_K^{n-1}$ is determined according to Table 1.*

PROOF: For full details of the proof see §§7,8 of [13]. □

5 Proof of Theorem 4.2

We need a few lemmas before we proceed with the proof of Theorem 4.2.

Lemma 5.1. *Let E/K be an elliptic curve with additive reduction. Assume that $\gcd(d, \text{char } k) = 1$. Then the group $E^0(K)$ is divisible by d .*

Table 1:

	c_p	$n = 2$	$n = 3$	$n = 4$
I_{2m}	2	$m + 1$, if $P \in E^0(K)$ 1, if $P \notin E^0(K)$	$m + 1$	$(m + 1)(m + 2)/2$, if $P \in E^0(K)$ $m + 1$, if $P \notin E^0(K)$
I_{2m}	$2m$	$m + 1$ if $\psi_E(P) \in 2\Phi_K(E)$ m if $\psi_E(P) \notin 2\Phi_K(E)$	$(m + 1)(2m + 1)/3$ if $3 \nmid m$ $m(2m + 3)/3 + 1$ if $3 \mid \gcd(m, \psi_E(P))$ $m(2m + 3)/3$ if $3 \mid m$ and $3 \nmid \psi_E(P)$	$m(m + 1)(m + 2)/3$ if $2 \nmid \psi_E(P)$ $m(m + 1)(m + 2)/3 + 1$ if $2 \mid \psi_E(P)$ $m(m + 1)(m + 2)/3 + 2$ if $4 \mid \psi_E(P)$
I_{2m+1}	1	$m + 1$	$m + 1$	$(m + 1)(m + 2)/2$
I_{2m+1}	$2m + 1$	$m + 1$	$(m + 1)(2m + 3)/3$ if $3 \nmid 2m + 1$ $(m + 2)(2m + 1)/3 + 1$ if $3 \mid \gcd(2m + 1, \psi_E(P))$ $(m + 2)(2m + 1)/3$ if $3 \mid 2m + 1$ and $3 \nmid \psi_E(P)$	$(m + 1)(m + 2)(2m + 3)/6$
II	1	1	1	1
III	1	1 if $\psi_E(P) = 1$, 2 if $\psi_E(P) = 0$	2	2 if $\psi_E(P) = 1$, 3 if $\psi_E(P) = 0$
IV	1 3	2	2 4 if $\psi_E(P) = 0$, 3 otherwise	3 5
I_0^*	1	2	3	4
I_{2m}^*	2	$m + 3$ if $P \in E^0(K)$ $m + 2$ if $P \notin E^0(K)$	$2m + 4$	$(m + 2)(m + 4)$ if $P \in E^0(K)$ $(m + 2)(m + 3)$ if $P \notin E^0(K)$
I_{2m}^*	4	$m + 5$ if $\psi_E(P) = (0, 0)$ $m + 2$ if $\psi_E(P) = (1, 1)$ 2 otherwise	$2m + 6$	$(m + 4)^2$ if $\psi_E(P) = (0, 0)$ $(m + 2)(m + 5)$ if $\psi_E(P) = (1, 1)$ $4m + 10$ otherwise
I_{2m+1}^*	2	$m + 4$ if $P \in E^0(K)$ $m + 2$ if $P \notin E^0(K)$	$2m + 5$	$(m + 3)(m + 4)$ if $P \in E^0(K)$ $(m + 2)(m + 4)$ if $P \notin E^0(K)$
I_{2m+1}^*	4	$m + 4$ if $2 \mid \psi_E(P)$ 2 otherwise	$2m + 7$	$(m + 3)(m + 6)$ if $\psi_E(P) = (0, 0)$ $(m + 4)^2$ if $\psi_E(P) = 2$ $4m + 12$ otherwise
IV*	1 3	3	4 8 if $\psi_E(P) = 0$, 6 otherwise	8 14
III*	2	4 if $\psi_E(P) = 0$, 2 otherwise	6	15 if $\psi_E(P) = 0$, 10 otherwise
II*	1	3	5	10

PROOF: Recall that $E^1(K) = \{P \in E(K) : \tilde{P} = \tilde{0}_E\}$. The group $E^0(K)/E^1(K)$ is isomorphic to k^+ because E has additive reduction. In particular, $E^0(K)/E^1(K)$ is divisible by d because $(d, \text{char } k) = 1$. But by the theory of formal groups, the group $E^1(K)$ is uniquely divisible by d . Therefore, $E^0(K)$ is divisible by d . $\square$

If E/K has non-split reduction type, then there exists a smallest finite unramified extension L/K of degree d over which E has split reduction. In fact, $d = 2$ except possibly when E has non-split reduction of type I_0^* then $d \in \{2, 3\}$. We define the norm map $\text{Norm}_{L/K}$ to be $\text{Norm}_{L/K} : E(L) \rightarrow E(K); Q \mapsto \sum_{i=1}^d Q^{\sigma^i}$, where $\text{Gal}(L/K) = \langle \sigma \rangle$.

Lemma 5.2. *Let L be the smallest unramified field over which E has split reduction. Then $\text{Norm}_{L/K} : E^0(L) \rightarrow E^0(K)$ is surjective.*

PROOF: We first treat the additive case. So assume that E has non-split additive reduction. Let $Q \in E^0(K)$. Since $E^0(K)$ is divisible by $d = [L : K]$, see Lemma 5.1, there is $Q' \in E^0(K)$ with $dQ' = Q$. Now we have $\text{Norm}_{L/K}(Q') = dQ' = Q$.

Now assume E has non-split multiplicative reduction. We always have $[L : K] = 2$. The non-singular reduction of E will be denoted by $\tilde{E}_{ns}$. Let ℓ be the residue field of L . We denote the image of σ under the isomorphism $\text{Gal}(L/K) \cong \text{Gal}(\ell/k)$ by σ again. Let $\text{Norm}_{\ell/k} : \tilde{E}_{ns}(\ell) \rightarrow \tilde{E}_{ns}(k); Q \mapsto Q + Q^\sigma$. Consider the following diagram.

$$\begin{array}{ccccccc} 0 & \longrightarrow & E^1(L) & \longrightarrow & E^0(L) & \longrightarrow & \tilde{E}_{ns}(\ell) \longrightarrow 0 \\ & & \downarrow \text{Norm}_{L/K} & & \downarrow \text{Norm}_{L/K} & & \downarrow \text{Norm}_{\ell/k} \\ 0 & \longrightarrow & E^1(K) & \longrightarrow & E^0(K) & \longrightarrow & \tilde{E}_{ns}(k) \longrightarrow 0 \end{array}$$

To prove that $\text{Norm}_{L/K} : E^0(L) \rightarrow E^0(K)$ is surjective, we only need to show the surjectivity of both $\text{Norm}_{L/K} : E^1(L) \rightarrow E^1(K)$ and $\text{Norm}_{\ell/k} : \tilde{E}_{ns}(\ell) \rightarrow \tilde{E}_{ns}(k)$. Let $Q \in E^1(K)$. Since $E^1(K)$ is divisible by 2, there is a $Q' \in E^1(K)$ such that $2Q' = Q$. So, $\text{Norm}_{L/K}(Q') = 2Q' = Q$.

Now we will show the surjectivity of $\text{Norm}_{\ell/k}$. The isomorphism $f : \tilde{E}_{ns}(\ell) \cong \ell^*$ induces an isomorphism $\tilde{E}_{ns}(k) \cong U := \{u \in \ell^* : \text{Norm}_{\ell/k}(u) = 1\}$. Moreover, the ℓ -automorphism $\sigma^{-1}f\sigma f^{-1} : \ell^* \rightarrow \ell^*$ is $u \mapsto u^{-1}$, see ([9], Exercise 10.2.7). Therefore, the map $\text{Norm}_{\ell/k}$ induces the map $\ell^* \rightarrow U; v \mapsto v/v^\sigma$. Now according to Hilbert's Theorem 90, for $u \in \ell^*$ we have $\text{Norm}_{\ell/k}(u) = 1$ if and only if $u = v/v^\sigma$ for some $v \in \ell^*$. Therefore, $\text{Norm}_{\ell/k}$ is surjective. $\square$

Lemma 5.3. *Let E/K be an elliptic curve with minimal proper regular model $E^{\min}$. Let $P \in E(K)$ and $(\Gamma_1, \dots, \Gamma_l)$ be an element of $S^n(P)$ endowed with the reduced structure. Then there exists a divisor D on $E^{\min} \rightarrow \text{Spec } \mathcal{O}_{K^{un}}$ such that:*

- (i) $(D|_E)^\sigma = D|_E$ for every $\sigma \in \text{Gal}(K^{un}/K)$.

- (ii) $D \cdot \Gamma_i = d_i$, where d_i is the number of times Γ_i appears in $(\Gamma_1, \dots, \Gamma_l)$.
- (iii) $D|_E \sim (n-1) \cdot 0 + P$.

PROOF: Assume first that E has reduction of type $I_m, m \geq 0$. Then E_k^{min} contains only multiplicity-1 irreducible components and we have $l = n$. Let $\ell = k(\Gamma_1, \dots, \Gamma_n)$. Let L/K be the unramified extension with residue field ℓ/k and $\text{Gal}(L/K) = \langle \sigma \rangle$. We have $[L : K] = [\ell : k] \leq 2$. Now we pick a point above each Γ_i . If Γ_i is defined over k , then choose $Q_i \in E(K)$ above Γ_i . If Γ_i is defined over ℓ , then pick $Q_i \in E(L)$ lying above Γ_i . Since $(\Gamma_1, \dots, \Gamma_n)$ is $\mathcal{G}_k$ -invariant, then $\Gamma_i^\sigma = \Gamma_j$ for some $j \in \{1, \dots, n\}$. We pick $Q_j := Q_i^\sigma$ to be the point above Γ_j . Since $\sum_{i=1}^n \delta_1(\Gamma_i) = \psi_E(P)$, it follows that $P' := \sum_{i=1}^n Q_i - P \in E^0(K)$. If there exists i such that $Q_i \in E(K)$, then replace Q_i with $Q_i - P'$. Otherwise, there exists i, j such that $Q_i \in E(L) \setminus E(K)$ and $Q_j = Q_i^\sigma$. According to Lemma 5.2, there is $Q' \in E^0(L)$ with $\text{Norm}_{L/K}(Q') = P'$. Now replace Q_i with $Q_i - Q'$, and Q_j with $Q_j - Q'^\sigma$. We set our divisor to be $D = \sum_{i=1}^n \overline{\{Q_i\}}$.

Now assume that E has additive reduction. If Γ_i is a multiplicity-1 component, then we pick Q_i above Γ_i as in the multiplicative case. We note that for non-split I_0^* , Γ_i might be defined over a cubic extension of k . In case Γ_i is defined over k and of multiplicity $m_i \geq 2$, then we choose $Q_i \in E$ above Γ_i , where $K(Q_i)/K$ is a totally ramified extension with $[K(Q_i) : K] = m_i$. In case of non-split reduction of type IV^* and $n = 4$, we can have $l = 2$ where Γ_1 is of multiplicity-2 and defined over a quadratic extension ℓ/k . Then we pick Q_1 above Γ_1 where $[L(Q_1) : L] = 2$ and we choose $Q_2 := Q_1^\sigma$. Let $G(Q_i) = \sum_{j=1}^{m_i} Q_i^{\sigma_j}$ where $\text{Gal}(K(Q_i)^{un}/K^{un}) = \langle \sigma_i \rangle$. Since $\sum_{i=1}^l \delta_{m_i}(\Gamma_i) = \psi_E(P)$, it follows from the definition of δ_{m_i} that $P' := \sum_i G(Q_i) - P \in E^0(K)$. By virtue of Lemma 5.1, there is $Q' \in E^0(K)$ with $nQ' = P'$. We set $Q'_i = Q_i - Q'$ for every i , and $D = \sum_i \overline{\{Q'_i\}}$. $\square$

PROOF OF THEOREM 4.2: We need to prove that the well-defined map

$$\lambda_n : \{ \text{minimal degree-}n\text{-models for } E \rightarrow \mathbb{P}_K^{n-1} \text{ up to isomorphism} \} \longrightarrow S^n(P)$$

is bijective. To prove that λ_n is injective, let $(\mathcal{C}_1, \alpha_1)$ and $(\mathcal{C}_2, \alpha_2)$ be two minimal degree- n -models for $E \rightarrow \mathbb{P}_K^{n-1}$. Assume that $\lambda_n(\mathcal{C}_1) = \lambda_n(\mathcal{C}_2)$. We want to show that $(\mathcal{C}_1, \alpha_1)$ and $(\mathcal{C}_2, \alpha_2)$ are isomorphic. The fact that they have the same corresponding tuples implies that they have the same strict transform in E_k^{min} , therefore $(\mathcal{C}_1, \alpha_1)$ and $(\mathcal{C}_2, \alpha_2)$ are isomorphic, see Theorem 2.1.

Now we will prove that λ_n is surjective. So assume that $(\Gamma_1, \dots, \Gamma_l) \in S^n(P)$ and we want to construct a minimal degree- n -model $(\mathcal{C}, \alpha)$ whose image under λ_n is $(\Gamma_1, \dots, \Gamma_l)$. By virtue of Lemma 5.3, there exists a divisor D on E^{min} such that $D|_E$ is K -rational, D intersects Γ_i as many times as its occurrence in the tuple, and $D|_E \sim (n-1) \cdot 0 + P$. Consider the following S -model for E

$$\mathcal{C} := \text{Proj} \left(\bigoplus_{m=0}^{\infty} H^0(E^{min}, \mathcal{O}_{E^{min}}(mD)) \right).$$

The model $\mathcal{C}$ is a minimal degree- n -model for $E \rightarrow \mathbb{P}_K^{n-1}$, see [12]. The special fiber $\mathcal{C}_k$ consists of the components in $(\Gamma_1, \dots, \Gamma_l)$ because D intersects these components in E_k^{min} and intersects no other components. Moreover, each component of $\mathcal{C}_k$ has degree equal to the number of iterations of its strict transform in $(\Gamma_1, \dots, \Gamma_l)$. We obtain α from the linear equivalence of $D|_E$ and $(n-1).0_E + P$. $\square$

6 Counting minimal global models

In this section we will be interested in attacking the global question. Let K be a number field of class number 1 with ring of integers $\mathcal{O}_K$. If $\mathfrak{p}$ is a prime in K , then we will denote the completion of K at $\mathfrak{p}$ by $K_{\mathfrak{p}}$, and its ring of integers by $\mathcal{O}_{K_{\mathfrak{p}}}$. Let C be a smooth genus one curve defined by an integral genus one equation of degree n over K . We assume moreover that $C(K_{\mathfrak{p}}) \neq \emptyset$ for every prime $\mathfrak{p}$ of K . A *minimal global degree- n -model* $(\mathcal{C}, \alpha)$ for $C \rightarrow \mathbb{P}_K^{n-1}$ consists of an $\text{Spec } \mathcal{O}_K$ -scheme defined by an $\mathcal{O}_K$ -integral genus one equation of degree n which is $\mathcal{O}_{K_{\mathfrak{p}}}$ -minimal at every prime $\mathfrak{p}$, and an isomorphism $\alpha : \mathcal{C}_K \cong C$ defined by an element in $\mathcal{G}_n(K)$. An *isomorphism* $\beta : (\mathcal{C}_1, \alpha_1) \cong (\mathcal{C}_2, \alpha_2)$ of degree- n -models for $C \rightarrow \mathbb{P}_K^{n-1}$ is an isomorphism $\beta : \mathcal{C}_1 \cong \mathcal{C}_2$ of $\text{Spec } \mathcal{O}_K$ -schemes defined by an element in $\mathcal{G}_n(\mathcal{O}_K)$ with $\beta_K = \alpha_2^{-1} \alpha_1$.

It is known that a minimal global degree-1-model for C/K , i.e., a Weierstrass model, exists and is unique up to isomorphism. A proof of the existence of a minimal global degree- n -model for C/K , where $C(K_{\mathfrak{p}}) \neq \emptyset$ for every $\mathfrak{p}$, when $n \in \{2, 3, 4\}$ can be found in [5]. The author gave a new proof in [12].

The problem of counting minimal global degree- n -models up to $\mathcal{O}_K$ -isomorphism can be tackled locally. We consider C as a curve over $K_{\mathfrak{p}}$ and use the previous section to determine the number $N_{\mathfrak{p}}$ of minimal degree- n -models for $C \rightarrow \mathbb{P}_{K_{\mathfrak{p}}}^{n-1}$ up to $\mathcal{O}_{K_{\mathfrak{p}}}$ -isomorphism. It turns out that we need only to investigate a finite set of primes. Then we collect the local data at the finite places using strong approximation.

If $m \in \mathcal{O}_K$, then we set $P(m) = \{\mathfrak{p} : \mathfrak{p} \text{ is prime, } \mathfrak{p}^2 \mid m\}$. The main result of this section is the following Theorem.

Theorem 6.1. *Let K be a number field of class number 1. Let $C \rightarrow \mathbb{P}_K^{n-1}$ be a smooth curve defined by a genus one equation of degree n , $n = 2, 3, 4$. Assume that $C(K_{\mathfrak{p}}) \neq \emptyset$ for every prime $\mathfrak{p}$. Let E/K be the Jacobian elliptic curve of C with minimal discriminant Δ . Let N and $N_{\mathfrak{p}}$ denote the numbers of minimal global degree- n -models for $C \rightarrow \mathbb{P}_K^{n-1}$, up to $\mathcal{O}_K$ -isomorphism, and minimal degree- n -models for $C \rightarrow \mathbb{P}_{K_{\mathfrak{p}}}^{n-1}$, up to $\mathcal{O}_{K_{\mathfrak{p}}}$ -isomorphism, respectively. Then*

$$N = \prod_{\mathfrak{p} \in P(\Delta)} N_{\mathfrak{p}}.$$

Theorem 6.1 follows immediately from the bijectivity of the following map

$$\begin{aligned} \{\text{minimal global degree-}n\text{-models for } C/K\} &\longrightarrow \prod_{\mathfrak{p} \in P(\Delta)} \{\text{minimal degree-}n\text{-models for } C/K_{\mathfrak{p}}\} \\ (\mathcal{C}, \alpha) &\mapsto ((\mathcal{C}, \alpha), \dots, (\mathcal{C}, \alpha)). \end{aligned}$$

Notice that the above two sets of degree- n -models are defined up to isomorphism. Furthermore, our work always enables us to compute $N_{\mathfrak{p}}$ when $\mathfrak{p}$ lies above a prime $p \geq 5$. Before proceeding with the proof of Theorem 6.1 we need the following Lemmas.

Lemma 6.2. *Let $A \in \mathrm{GL}_n(K_{\mathfrak{p}}) \cap \mathrm{Mat}_n(\mathcal{O}_{K_{\mathfrak{p}}})$ have coprime entries. Assume that $\mathfrak{p} = \pi\mathcal{O}_K$. Then there exist matrices $U \in \mathrm{GL}_n(\mathcal{O}_K)$ and $V \in \mathrm{GL}_n(\mathcal{O}_{K_{\mathfrak{p}}})$ such that $A = VDU$, where $D = \mathrm{diag}(\pi^{r_1}, \dots, \pi^{r_{n-1}}, 1)$ and $r_1 \geq \dots \geq r_{n-1}$.*

PROOF: We claim that there exists a matrix $B \in \mathrm{GL}_n(K) \cap \mathrm{Mat}_n(\mathcal{O}_K)$ such that $V' := BA^{-1} \in \mathrm{GL}_n(\mathcal{O}_{K_{\mathfrak{p}}})$. Granted this claim we write the Smith Normal Form for the matrix B , so we have $B = U'D'DU$ where $U, U' \in \mathrm{GL}_n(\mathcal{O}_K)$, D' is a diagonal matrix whose entries are not divisible by π , and $D = \mathrm{diag}(\pi^{r_1}, \dots, \pi^{r_{n-1}}, 1)$, $r_1 \geq \dots \geq r_{n-1}$. Then we set $V := V'^{-1}U'D' \in \mathrm{GL}_n(\mathcal{O}_{K_{\mathfrak{p}}})$, hence we are done.

To prove the claim, assume that $A = (a_{ij})_{i,j}$. Recall that every element in $\mathcal{O}_{K_{\mathfrak{p}}}$ can be written uniquely as $\sum_{i \geq 0} a_i \pi^i$, where $a_i \in \mathcal{O}_K$ lies in a set of representatives of $\mathcal{O}_K/\mathfrak{p}$. By a continuity argument, there exists an integer $m > 0$ such that the matrix $B = (a_{ij} \bmod \pi^m)_{i,j} \in \mathrm{GL}_n(K) \cap \mathrm{Mat}_n(\mathcal{O}_K)$. Now we have $BA^{-1} \equiv \mathrm{id}_n \bmod \pi^m$ and hence $BA^{-1} \in \mathrm{GL}_n(\mathcal{O}_{K_{\mathfrak{p}}})$. $\square$

Lemma 6.3. *Let ϕ and ϕ' be two minimal $\mathcal{G}_n(K_{\mathfrak{p}})$ -equivalent genus one equations of degree n with coefficients in $\mathcal{O}_K$ and $\mathcal{O}_{K_{\mathfrak{p}}}$ respectively. Then ϕ' is $\mathcal{G}_n(\mathcal{O}_{K_{\mathfrak{p}}})$ -equivalent to a minimal genus one equation of degree n whose coefficients lie in $\mathcal{O}_K$.*

PROOF: Assume that ϕ' is obtained from ϕ via $[\mu_n, A_n]$ in $\mathcal{G}_n(K_{\mathfrak{p}})$. For $r \in K_{\mathfrak{p}}^*$, the following transformations are identical:

$$[\mu_n, A_n] = [r^{-2}\mu_n, rA_n] \text{ when } n = 2, 4, \text{ and } [\mu_3, A_3] = [r^{-3}\mu_3, rA_3].$$

Therefore, we can assume that A_n has coprime entries in $\mathcal{O}_{K_{\mathfrak{p}}}$. Lemma 6.2 allows us to write $A_n = V_n D_n U_n$ where $V_n \in \mathrm{GL}_n(\mathcal{O}_{K_{\mathfrak{p}}})$, $U_n \in \mathrm{GL}_n(\mathcal{O}_K)$, and $D_n = \mathrm{diag}(\pi^{r_1}, \dots, \pi^{r_{n-1}}, 1)$ where $\mathfrak{p} = \pi\mathcal{O}_K$. Similarly, we can write $\mu_4 = \nu'_4 \tau_4 \nu_4$ where $\nu'_4 \in \mathrm{GL}_2(\mathcal{O}_{K_{\mathfrak{p}}})$, $\nu_4 \in \mathrm{GL}_2(\mathcal{O}_K)$ and $\tau_4 = \mathrm{diag}(\pi^{-m_1}, \pi^{-m_2})$.

Let ψ be the $\mathcal{O}_K$ -integral genus one equation obtained from ϕ via the transformation $[1, U_n]$ when $n = 2, 3$, and via $[\nu_4, U_4]$ when $n = 4$. Then ψ lies in the same $\mathcal{G}_n(\mathcal{O}_K)$ -equivalence class as ϕ . Let ϕ'' be the genus one equation obtained from ψ via the transformation $[\mu'_n, D_n]$, where $\mu'_2 = (\det D_2)^{-2}$, $\mu'_3 = (\det D_3)^{-1}$ and $\mu'_4 = \tau_4$. It is clear that ϕ'' is $\mathcal{G}_n(\mathcal{O}_{K_{\mathfrak{p}}})$ -equivalent to ϕ' . We claim that ϕ'' has coefficients in $\mathcal{O}_K$. If it is not the case, then some of the coefficients

of the polynomials defining ϕ'' would lie in $\frac{1}{\pi^r}\mathcal{O}_K$ for some $r > 0$. But ϕ'' is obtained from ϕ' via $[\omega_n, V_n^{-1}]$, where $\omega_n \in \mathcal{O}_{K_{\mathfrak{p}}}^*$ when $n = 2, 3$, and $\omega_4 \in \text{GL}_2(\mathcal{O}_{K_{\mathfrak{p}}})$. Since ϕ' is $\mathcal{O}_{K_{\mathfrak{p}}}$ -integral, it follows that ϕ'' should be $\mathcal{O}_{K_{\mathfrak{p}}}$ -integral, which is a contradiction. $\square$

The following lemma, ([14], Chapter IV, Lemma 9.5), will be used to justify our choice of the set of prime numbers $P(\Delta)$.

Lemma 6.4. *Let $E/K_{\mathfrak{p}}$ be an elliptic curve with discriminant Δ . If $\nu_{\mathfrak{p}}(\Delta) = 1$, then E has reduction of type I_1 .*

PROOF OF THEOREM 6.1: First we will show that the map λ is well defined. Let $(\mathcal{C}_1, \alpha_1)$ and $(\mathcal{C}_2, \alpha_2)$ be two isomorphic minimal global degree- n -models for $C \rightarrow \mathbb{P}_K^{n-1}$. Then $\alpha := \alpha_2^{-1}\alpha_1 : (\mathcal{C}_1)_{K_{\mathfrak{p}}} \rightarrow (\mathcal{C}_2)_{K_{\mathfrak{p}}}$ is defined by an element in $\mathcal{G}_n(\mathcal{O}_K) \hookrightarrow \mathcal{G}_n(\mathcal{O}_{K_{\mathfrak{p}}})$ for every prime $\mathfrak{p}$, i.e., $(\mathcal{C}_1, \alpha_1)$ and $(\mathcal{C}_2, \alpha_2)$ have the same image under λ .

To show that λ is injective, let $(\mathcal{C}_1, \alpha_1)$ and $(\mathcal{C}_2, \alpha_2)$ be two minimal global degree- n -models for $C \rightarrow \mathbb{P}_K^{n-1}$ with the same image under λ . We need to show that $(\mathcal{C}_1, \alpha_1)$ and $(\mathcal{C}_2, \alpha_2)$ are isomorphic. Let $\alpha := \alpha_2^{-1}\alpha_1$. The map α is defined by an element $[\mu, A] \in \mathcal{G}_n(K)$. We can assume that $A \in \text{Mat}_n(\mathcal{O}_K)$ has coprime entries. Since $(\mathcal{C}_1, \alpha_1)$ and $(\mathcal{C}_2, \alpha_2)$ have the same image under λ , in particular they are $\mathcal{O}_{K_{\mathfrak{p}}}$ -isomorphic for every $\mathfrak{p} \in P(\Delta)$, it follows that $A \in \text{GL}_n(\mathcal{O}_{K_{\mathfrak{p}}})$, and so $\mathfrak{p} \nmid \det A$. If $\mathfrak{p} \notin P(\Delta)$, then $E/K_{\mathfrak{p}}$ has either reduction types I_0 or I_1 , see Lemma 6.4. But according to Corollary 2.2, when E has either reduction types I_0 or I_1 , there is a unique degree- n -model for $C \rightarrow \mathbb{P}_{K_{\mathfrak{p}}}^{n-1}$. That means that for $\mathfrak{p} \notin P(\Delta)$, $(\mathcal{C}_1, \alpha_1)$ and $(\mathcal{C}_2, \alpha_2)$ are isomorphic as degree- n -models for $C \rightarrow \mathbb{P}_{K_{\mathfrak{p}}}^{n-1}$. Hence $A \in \text{GL}_n(\mathcal{O}_{K_{\mathfrak{p}}})$ for every prime $\mathfrak{p}$, in particular $\mathfrak{p} \nmid \det A$. Thus $\det A \in \mathcal{O}_K^*$ and $A \in \text{GL}_n(\mathcal{O}_K)$.

Now we will prove the surjectivity of λ . We will assume without loss of generality that the defining genus one equation ϕ of C has coefficients in $\mathcal{O}_K$ and that the associated discriminant is everywhere minimal.

If $P(\Delta) = \emptyset$, then the scheme defined by ϕ is the unique minimal global degree- n -model for $C \rightarrow \mathbb{P}_K^{n-1}$. So, assume $P(\Delta) = \{\mathfrak{p}_1, \dots, \mathfrak{p}_m\}$, $m \geq 1$. Let $(\mathcal{C}_i, \alpha_i)$, where $1 \leq i \leq m$, be a minimal degree- n -model for $C \rightarrow \mathbb{P}_{K_{\mathfrak{p}_i}}^{n-1}$. We want to construct a minimal global degree- n -model $(\mathcal{C}, \alpha)$ for $C \rightarrow \mathbb{P}_K^{n-1}$ such that $\alpha^{-1}\alpha_i : (\mathcal{C}_i)_K \rightarrow \mathcal{C}_K$ is defined by an element in $\mathcal{G}_n(\mathcal{O}_{K_{\mathfrak{p}_i}})$ for each i . Let ϕ_i be the defining genus one equation of $\mathcal{C}_i$. By virtue of Lemma 6.3, we can assume that ϕ_i has coefficients in $\mathcal{O}_K$ and is obtained from ϕ via $[\mu_i, D_i U_i]$ where $D_i = \text{diag}(\pi_i^{r_{i,1}}, \dots, \pi_i^{r_{i,n-1}}, 1)$, where $\mathfrak{p}_i = \pi_i \mathcal{O}_K$, $U_i \in \text{GL}_n(\mathcal{O}_K)$ and μ_i is a scaling element. In fact, we can assume that $U_i \in \text{SL}_n(\mathcal{O}_K)$ by acting by a diagonal matrix whose entries lie in $\mathcal{O}_K^*$. Given integers $m_i > 0$, there exists a matrix $U \in \text{SL}_n(\mathcal{O}_K)$ such that $U \equiv U_i \pmod{\pi_i^{m_i}}$ for every i , see ([6], Lemma 3.2). We note that $\prod_{i=1}^m D_i U U_i^{-1} D_i^{-1} \equiv \prod_{i \neq j} D_i \pmod{\pi_j^{m_j}}$. Therefore, the genus one equation ψ obtained from ϕ via the transformation $[\prod_{i=1}^m \mu_i, \prod_{i=1}^m D_i U]$ is $\mathcal{G}_n(\mathcal{O}_{K_{\mathfrak{p}_j}})$ -equivalent to ϕ_j for every j .

Now we want to show that ψ is $\mathcal{O}_K$ -integral. This will imply that ψ defines a minimal global degree- n -model for $C \rightarrow \mathbb{P}_K^{n-1}$ which is $\mathcal{O}_{K_{\mathfrak{p}_j}}$ -isomorphic to $(\mathcal{C}_j, \alpha_j)$ for every j . Hence we will

be done with the surjectivity. Assume on the contrary that ψ is not $\mathcal{O}_K$ -integral. Since ψ is obtained from the $\mathcal{O}_K$ -integral genus one equation ϕ via $[\prod_{i=1}^m \mu_i, \prod_{i=1}^m D_i U]$, it follows that some of the coefficients of the defining polynomials of ψ lie in $\frac{1}{b}\mathcal{O}_K$, $b = \pi_1^{l_1} \pi_2^{l_2} \dots \pi_m^{l_m}$, where $l_i \geq 0$ and $l_j > 0$ for some $j \in \{1, \dots, m\}$. We have shown that ψ is $\mathcal{G}_n(\mathcal{O}_{K_{p_j}})$ -equivalent to the $\mathcal{O}_K$ -integral genus one equation ϕ_j . It follows that ψ is $\mathcal{O}_{K_{p_j}}$ -integral, which is a contradiction. $\square$

The calculations included in the following examples are performed using MAGMA, see [3]. $N_n(T)$ will denote the number of minimal degree- n -models when the reduction of the Jacobian is of type T .

Example 6.5. Consider the elliptic curve $E : y^2 + xy = x^3 - x^2 - 617x + 5916/\mathbb{Q}$. It has reduction of types III^* and I_2 at its bad primes 5 and 19 respectively. The following minimal global genus one equation ϕ_3 of degree 3 defines an everywhere locally soluble element C in the 3-Selmer group of E .

$$\begin{aligned} 21686353648850x^3 &+ 234081254700017x^2y + 9338329782950x^2z + 842219868972245xy^2 + \\ 67198263238095xyz &+ 1340388284750xz^2 + 1010096983050575y^3 + 120889031707155y^2z + \\ 4822691362750yz^2 &+ 64131409475z^3 = 0. \end{aligned}$$

The number of minimal global degree-3-models for $C \rightarrow \mathbb{P}_K^2$ is $N_3(\text{III}^*) \times N_3(\text{I}_2) = 6 \times 2 = 12$, see Theorem 6.1. These models have defining genus one equations obtained from ϕ_3 via the following transformations in $\mathcal{G}_3(\mathbb{Q})$.

$$\begin{aligned} [1, \text{id}_3], & \quad [1/5, \text{diag}(5, 1, 1)], & [1/5, \text{diag}(1, 5, 1)], & \quad [1/25, \text{diag}(5, 5, 1)], \\ [1/25, \text{diag}(5, 1, 5)], & [1/25, \text{diag}(1, 25, 1)], & [1/19, \text{diag}(1, 1, 19)], & [1/95, \text{diag}(5, 1, 19)], \\ [1/95, \text{diag}(1, 5, 19)], & [1/475, \text{diag}(5, 5, 19)], & [1/475, \text{diag}(5, 1, 95)], & [1/475, \text{diag}(1, 25, 19)]. \end{aligned}$$

We notice that the genus one equations ϕ_3 is not reduced in the sense of [5], but we preferred to write it in that way to make the transformations as simple as possible.

Example 6.6. Let $E : y^2 + xy + y = x^3 - 4x - 3/\mathbb{Q}$. The curve $C : y^2 = -3x^4 + 2x^3 + 7x^2 - 2x - 3$ represents an element in the 2-Selmer group of E . A second 2-descent on C gives the following minimal global genus one equation ϕ_4 of degree 4.

$$\begin{aligned} x_1^2 - x_1x_3 - x_2^2 + x_2x_4 + x_3^2 &= 0, \\ x_1x_4 + x_2^2 + x_2x_3 - x_2x_4 + x_3^2 - x_3x_4 &= 0. \end{aligned}$$

The equation ϕ_4 defines a smooth genus one curve $C_4/\mathbb{Q}$. The discriminant Δ of E is 185. Therefore, we have $P(\Delta) = \emptyset$. Hence the minimal global degree-4-model $\mathcal{C}$ defined by ϕ_4 is unique, see Theorem 6.1.

Acknowledgements. This paper is based on the author's Ph.D. thesis [13] at Cambridge University. The author would like to thank his supervisor Dr. Tom Fisher for all the encouragement and guidance through this work.

References

- [1] S.Y. An, S.Y. Kim, D.C. Marshall, S.H. Marshall, W.G. McCallum, and A.R. Perlis. Jacobians of genus one curves. *J. Number Theory*, 90(2):304–315, 2001.
- [2] B.J. Birch and H.P.F. Swinnerton-Dyer. Notes on elliptic curves I. *J. reine angew. Math.*, 212:7–25, 1963.
- [3] W. Bosma, J. Cannon, and C. Playoust. The Magma algebra system I: The user language. *J. Symb. Comb.*, 24:235–265, 1997.
- [4] T. Bromwich. *Quadratic forms and their classification by means of invariant factors*. Cambridge Tracts in Mathematics and Mathematical Physics, 1906.
- [5] J.E. Cremona, T.A. Fisher, and M. Stoll. Minimisation and reduction of 2-, 3- and 4-coverings of elliptic curves, preprint.
- [6] T.A. Fisher. A new approach to minimising binary quartics and ternary cubics. *Math. Res. Lett.*, 14:597–613, 2007.
- [7] T.A. Fisher. The invariants of a genus one curve. *Proc. Lond. Math. Soc.*, 97(3):753–782, 2008.
- [8] Q. Liu. Modèles entiers des courbes hyperelliptiques sur un corps de valuations discrète. *Trans. Amer. Math. Soc.*, 348(11):4577–4610, November 1996.
- [9] Q. Liu. *Algebraic Geometry and Arithmetic Curves*, volume 6 of *Oxford Graduate Texts in Mathematics*. Oxford University Press, Oxford, 2002.
- [10] D. Lorenzini. Models of curves and wild ramification, preprint.
- [11] B. Poonen. An explicit algebraic family of genus-one curves violating the Hasse principle. *J. Théor. Nombres Bordeaux*, 13(1):263–274, 2001.
- [12] M. Sadek. Minimal genus one curves, preprint.
- [13] M. Sadek. *Models of genus one curves*. PhD thesis, Cambridge University, 2009.
- [14] J. Silverman. *Advanced topics in the arithmetic of elliptic curves*. GTM 151. Springer-Verlag, 1995.
- [15] M. Stoll and J.E. Cremona. Minimal models for 2-coverings of elliptic curves. *LMS J. Comput. Math.*, 5:220–243, 2002.
- [16] T.O. Womack. *Explicit descent on elliptic curves*. PhD thesis, University of Nottingham, 2003.